


REMARKS

Applicants request entry of this amendment in adherence with 37 C.F.R. §§1.821 to 1.825. If the Examiner believes a telephone conference would expedite prosecution of this application, please telephone the undersigned at 415-576-0200.

Respectfully submitted,



Jean M. Lockyer
Reg. No. 44,879

TOWNSEND and TOWNSEND and CREW LLP
Two Embarcadero Center, 8th Floor
San Francisco, California 94111-3834
Tel: (415) 576-0200
Fax: (415) 576-0300
JML:dmw

10021660-120601

VERSION WITH MARKINGS TO SHOW CHANGES MADE

In the Specification:

Paragraph beginning at line 23 of page 42 has been amended as follows:

Various tag polypeptides and their respective antibodies are well known in the art. Examples include poly-histidine (poly-his) or poly-histidine-glycine (poly-his-gly) tags; HIS6 (SEQ ID NO:135) and metal chelation tags, the flu HA tag polypeptide and its antibody 12CA5 [Field *et al.*, *Mol. Cell. Biol.*, 8:2159-2165 (1988)]; the c-myc tag and the 8F9, 3C7, 6E10, G4, B7 and 9E10 antibodies thereto [Evan *et al.*, *Molecular and Cellular Biology*, 5:3610-3616 (1985)]; and the Herpes Simplex virus glycoprotein D (gD) tag and its antibody [Paborsky *et al.*, *Protein Engineering*, 3(6):547-553 (1990)]. Other tag polypeptides include the Flag-peptide [Hopp *et al.*, *BioTechnology*, 6:1204-1210 (1988)]; the KT3 epitope peptide [Martin *et al.*, *Science*, 255:192-194 (1992)]; tubulin epitope peptide [Skinner *et al.*, *J. Biol. Chem.*, 266:15163-15166 (1991)]; and the T7 gene 10 protein peptide tag [Lutz-Freyermuth *et al.*, *Proc. Natl. Acad. Sci. USA*, 87:6393-6397 (1990)].

Paragraph beginning at line 3 of page 82 has been amended as follows:

AAA4 DNA sequence (SEQ ID NO:1):

Gene name: CGI-100 protein

Unigene number: Hs.275253

Probeset Accession #: AA089688

Nucleic Acid Accession #: NM_016040 cluster

Coding sequence: 142-831 (predicted start/stop codons underlined)

Paragraph beginning at line 32 of page 82 has been amended as follows:

AAA7 DNA sequence (SEQ ID NO:2):

Gene name: Endothelial differentiation, sphingolipid G-protein-coupled receptor, 1 (EDG1)

Unigene number: Hs.154210

Probeset Accession #: M31210

Nucleic Acid Accession #: NM_001400 cluster

Coding sequence: 251-1396 (predicted start/stop codons underlined)

Paragraph beginning at line 20 of page 83 has been amended as follows:

AAB3 DNA sequence (SEQ ID NO:3):

Gene name: Solute carrier family 20 (phosphate transporter), member 1, Human leukaemia virus receptor 1 (GLVR1)
Unigene number: Hs.78452
Probeset Accession #: L20859
Nucleic Acid Accession #: NM_005415 cluster
Coding sequence: predicted 371-2410 (predicted start/stop codons underlined)

Paragraph beginning at line 16 of page 84 has been amended as follows:

AAB4 DNA sequence (SEQ ID NO:4):

Gene name: Matrix metalloproteinase 10 (stromelysin 2)
Unigene number: Hs.2258
Probeset Accession #: X07820
Nucleic Acid Accession #: NM_002425
Coding sequence: predicted 23-1453 (predicted start/stop codons underlined)

Paragraph beginning at line 55 of page 84 has been amended as follows:

AAB6 DNA sequence (SEQ ID NO:5):

Gene name: Podocalyxin-like
Unigene number: Hs.16426
Probeset Accession #: U97519
Nucleic Acid Accession #: NM_005397 cluster
Coding sequence: 251-1837 (predicted start/stop codons underlined)

Paragraph beginning at line 26 of page 86 has been amended as follows:

AAB8 DNA sequence (SEQ ID NO:6):

Gene name: EGF-containing fibulin-like extracellular matrix protein 1
Unigene number: Hs.76224
Probeset Accession #: U03877
Nucleic Acid Accession #: NM_004105 Transcript variant 1
Coding sequence: 150-1631 (predicted start/stop codons underlined)

10021650.120601

Paragraph beginning at line 13 of page 87 has been amended as follows:

AAB9 DNA sequence (SEQ ID NO:7):

Gene name: Melanoma adhesion molecule, MUC 18 glycoprotein
Unigene number: Hs.211579
Probeset Accession #: M28882
Nucleic Acid Accession #: NM_006500 cluster
Coding sequence: 27-1967 (predicted start/stop codons underlined)

Paragraph beginning at line 14 of page 88 has been amended as follows:

AAC1 DNA sequence (SEQ ID NO:8):

Gene name: Matrix metalloproteinase 1 (interstitial collagenase)
Unigene number: Hs.83169
Probeset Accession #: X54925
Nucleic Acid Accession #: NM_002421 cluster
Coding sequence: 69-1478 (predicted start/stop codons underlined)

Paragraph beginning at line 56 of page 88 has been amended as follows:

AAC3 DNA sequence (SEQ ID NO:9):

Gene name: Branched chain aminotransferase 1, cytosolic
Unigene number: Hs.157205
Probeset Accession #: AA423987
Nucleic Acid Accession #: NM_005504 cluster
Coding sequence: 1-1155 (predicted start/stop codons underlined)

Paragraph beginning at line 17 of page 89 has been amended as follows:

ACG4 DNA sequence (SEQ ID NO:10):

Gene name: Pentaxin-related gene, rapidly induced by IL-1 beta
Unigene number: Hs.2050
Probeset Accession #: M31166
Nucleic Acid Accession #: NM_002852 cluster
Coding sequence: 68-1213 (predicted start/stop codons underlined)

10021660-120601

Paragraph beginning at line 57 of page 89 has been amended as follows:

ACK5 DNA sequence (SEQ ID NO:11):

Gene name: Von Willebrand factor; Coagulation factor VIII
Unigene number: Hs.110802
Probeset Accession #: M10321
Nucleic Acid Accession #: NM_000552
Coding sequence: 311-8752 (predicted start/stop codons underlined)

Paragraph beginning at line 11 of page 92 has been amended as follows:

AAC7 DNA sequence (SEQ ID NO:12):

Gene name: KIAA1294 protein
Probeset Accession #: AA432248
Nucleic Acid Accession #: AB037715
Coding sequence: 370-3489 (predicted start/stop codons underlined)

Paragraph beginning at line 65 of page 93 has been amended as follows:

ACG8 DNA sequence (SEQ ID NO:13):

Gene name: ubiquitin E3 ligase SMURF2
Unigene number: Hs.21806 (3'UTR only)
Probeset Accession #: AA398243
Nucleic Acid Accession #: AF301463 cluster
Coding sequence: 9-2255 (predicted start/stop codons underlined)

Paragraph beginning at line 44 of page 94 has been amended as follows:

ACH1 DNA sequence (SEQ ID NO:14):

Gene name: EST
Unigene number: Hs.30089
Probeset Accession #: AA410480
CAT cluster#: 96816_1
Coding sequence: Partial sequence, possible frameshift. Predicted stop codon underlined.

10021660-120601
T09021-099T2001

Paragraph beginning at line 2 of page 95 has been amended as follows:

ACJ2 DNA sequence (SEQ ID NO:15):

Gene name: Complement component C1q receptor

Unigene number: Hs.97199

Probeset Accession #: AA487558

Nucleic Acid Accession #: NM_012072

Coding sequence: 149-2107. Predicted start/stop codons underlined

Paragraph beginning at line 55 of page 96 has been amended as follows:

ACJ3 DNA sequence (SEQ ID NO:16):

Gene name: FLT1/vascular endothelial growth factor receptor

Unigene number: Hs.138671

Probeset Accession #: AA047437

Nucleic Acid Accession #: NM_002019

Coding sequence: 250-4266 (predicted start/stop codons underlined)

Paragraph beginning at line 56 of page 98 has been amended as follows:

ACJ9 DNA sequence (SEQ ID NO:17):

Gene name: Purine nucleoside phosphorylase

Unigene number: Hs.75514

Probeset Accession #: K02574

Nucleic acid Accession #: X00737 cluster

Coding sequence: 110-979 (predicted start/stop codons underlined)

Paragraph beginning at line 21 of page 99 has been amended as follows:

ACK4 DNA sequence (SEQ ID NOS:18 and 19):

Gene name: EST

Unigene number: Hs.265499

Probeset Accession #: R68763

CAT cluster#: Cluster 46668_2

Sequence: Both the EST corresponding to the probeset accession and exon prediction; number and the CAT cluster align with the Homo sapiens BAC clone AC009414 RP11-490M8. Using FGESH, 2 exons predicted on this BAC clone upstream of the probeset.

Predicted exon 1: bases 5808-5837 of BAC clone AC009414

Paragraph beginning at line 58 of page 99 has been amended as follows:

Predicted ACK4 gene seq (predicted start/stop codons underlined) (SEQ ID NO:19):

10021660-120601

Paragraph beginning at line 20 of page 100 has been amended as follows:

AAA8 DNA sequence (SEQ ID NO:20):

Gene name: ETL protein, with extended open reading frame

Unigene number: Hs.57958

Probeset Accession #: D58024

Nucleotide Accession #: AF192403

Coding sequence: 151-2136. Underlined sequences correspond to extended sequence not included in AF192403.

Paragraph beginning at line 11 of page 101 has been amended as follows:

AAC6 DNA sequence (SEQ ID NO:21):

Gene name: Homo sapiens cDNA FLJ13465 fis, clone PLACE1003493, weakly similar to endothelial cell multimerin precursor

Unigene number: Hs.134797

Probeset Accession #: AA025351

Nucleotide Accession #: AK023527

Coding sequence: predicted 75-2921

Extended sequence: 729-3465 (underlined sequence)

Paragraph beginning at line 18 of page 102 has been amended as follows:

ACH7 DNA sequence (SEQ ID NOS:22-24):

Gene name: ESTs

Unigene number: Hs.3807

Probeset Accession #: AA292694

BAC Accession #: AL161751

FGENESH predicted exons: FGENESH predicts 2 exons on the minus strand of AL161751 upstream of the ACH7 probeset.

Paragraph beginning at line 26 of page 102 has been amended as follows:

FGENESH predicted exon 1 (SEQ ID NO:22):

Paragraph beginning at line 32 of page 102 has been amended as follows:

FGENESH predicted exon 2 (SEQ ID NO:23):

10021660-120601
T09021-099T2001

Paragraph beginning at line 63 of page 102 has been amended as follows:

ACH7 predicted coding seq (predicted start/stop codons underlined) (SEQ ID NO:24):

Paragraph beginning at line 31 of page 103 has been amended as follows:

AAD3 DNA sequence (SEQ ID NO:25):

Gene name: ESTs

Unigene number: Hs.17404

Probeset Accession #: N39584

Nucleic Acid Accession #: N39584

Coding sequence: no identified ORF; possible frameshifts

Paragraph beginning at line 64 of page 103 has been amended as follows:

AAD4 DNA sequence (SEQ ID NO:26):

Gene name: ERG

Unigene number: Hs.279477 / Hs.45514

Probeset Accession #: R32894

Nucleic Acid Accession #: M17254

Coding sequence: 257-1645 (predicted start/stop codons underlined)

Paragraph beginning at line 58 of page 104 has been amended as follows:

AAD5 DNA sequence (SEQ ID NO:27):

Gene name: activin A receptor type II-like 1 (ALK-1)

Unigene number: Hs.8881 / Hs.172670

Probeset Accession #: T57112

Nucleic Acid Accession #: NM_000020

Coding sequence: 283-1794 (predicted start/stop codons underlined)

Paragraph beginning at line 3 of page 106 has been amended as follows:

AAD8 DNA sequence (SEQ ID NO:28):

Gene name: ESTs

Unigene number: Hs.144953

Probeset Accession #: AA404418

Nucleic Acid Accession #: n/a

Coding sequence: no ORF identified; possible frameshifts

10021660-120601

Paragraph beginning at line 39 of page 106 has been amended as follows:

ACA2 DNA sequence (SEQ ID NO:29):

Gene name: EST

Unigene number: Hs.16450

Probeset Accession #: AA478778

Nucleic Acid Accession #: AA478778

Coding sequence: no ORF identified; possible frameshifts

Paragraph beginning at line 1 of page 107 has been amended as follows:

ACA4 DNA sequence (SEQ ID NO:30):

Gene name: alpha satellite junction DNA sequence

Unigene number: Hs.247946

Probeset Accession #: M21305

Nucleic Acid Accession #: M21305

Coding sequence: 1-165 (predicted start/stop codons underlined)

Paragraph beginning at line 13 of page 107 has been amended as follows:

ACG6 DNA sequence (SEQ ID NO:31):

Gene name: intercellular adhesion molecule 2 (ICAM2)

Unigene number: Hs.83733

Probeset Accession #: M32334

Nucleic Acid Accession #: NM_000873

Coding sequence: 63-890 (predicted start/stop codons underlined)

Paragraph beginning at line 40 of page 107 has been amended as follows:

ACG7 DNA sequence (SEQ ID NO:32):

Gene name: Cadherin 5, VE-cadherin (CDH5)

Unigene number: Hs.76206

Probeset Accession #: X79981

Nucleic Acid Accession #: NM_001795

Coding sequence: 25-2379 (predicted start/stop codons underlined)

10021660-120601

Paragraph beginning at line 48 of page 108 has been amended as follows:

ACG9 DNA sequence (SEQ ID NO:33):
Gene name: lysyl oxidase-like 2 (LOXL2)
Unigene number: Hs.83354
Probeset Accession #: U89942
Nucleic Acid Accession #: NM_002318 cluster
Coding sequence: 248-2572 (predicted start/stop codons underlined)

Paragraph beginning at line 47 of page 109 has been amended as follows:

ACH2 DNA sequence (SEQ ID NO:34):
Gene name: TIE tyrosine-protein kinase
Unigene number: Hs.78824
Probeset Accession #: X60957
Nucleic Acid Accession #: NM_005424 cluster
Coding sequence: 37-3452 (predicted start/stop codons underlined)

Paragraph beginning at line 53 of page 110 has been amended as follows:

ACH3 DNA sequence (SEQ ID NO:35):
Gene name: placental growth factor (PGF; PlGF1; VEGF-related protein)
Unigene number: Hs.2894
Probeset Accession #: X54936
Nucleic Acid Accession #: NM_002632 cluster
Coding sequence: 322-768 (predicted start/stop codons underlined)

Paragraph beginning at line 22 of page 111 has been amended as follows:

ACH4 DNA sequence (SEQ ID NO:36):
Gene name: nidogen 2 (NID2)
Unigene number: Hs.82733
Probeset Accession #: D86425
Nucleic Acid Accession #: NM_007361 cluster
Coding sequence: 1-4131 (predicted start/stop codons underlined)

10021650-120601
T0909T-0992200F

Paragraph beginning at line 44 of page 112 has been amended as follows:

ACH5 DNA sequence (SEQ ID NO:37):

Gene name: SNL (singled-like; sea urchin fascin homolog-like)
Unigene number: Hs.118400
Probeset Accession #: U03057
Nucleic Acid Accession #: NM_003088
Coding sequence: 112-1593 (predicted start/stop codons underlined)

Paragraph beginning at line 32 of page 113 has been amended as follows:

ACH6 DNA sequence (SEQ ID NO:38):

Gene name: endothelial protein C receptor (EPCR; PROCR)
Unigene number: Hs.82353
Probeset Accession #: L35545
Nucleic Acid Accession #: NM_006404
Coding sequence: 25-741 (predicted start/stop codons underlined)

Paragraph beginning at line 63 of page 113 has been amended as follows:

ACH8 DNA sequence (SEQ ID NO:39):

Gene name: melanoma adhesion molecule (MCAM; MUC18)
Unigene number: Hs.211579
Probeset Accession #: D51069
Nucleic Acid Accession #: NM_006500
Coding sequence: 27-1967 (predicted start and stop codons underlined)

Paragraph beginning at line 64 of page 114 has been amended as follows:

ACH9 DNA sequence (SEQ ID NO:40):

Gene name: endothelin-1 (EDN1)
Unigene number: Hs.2271
Probeset Accession #: J05008
Nucleic Acid Accession #: NM_001955
Coding sequence: 337-975 (predicted start/stop codons underlined)

10021660-120601
109021-09972001

Paragraph beginning at line 26 of page 115 has been amended as follows:

ACJ1 DNA sequence (SEQ ID NO:41):

Gene name: BMX non-receptor tyrosine kinase

Unigene number: Hs.27372

Probeset Accession #: X83107

Nucleic Acid Accession #: NM_001721

Coding sequence: 34-2061 (predicted start/stop codons underlined)

Paragraph beginning at line 8 of page 116 has been amended as follows:

ACJ4 DNA sequence (SEQ ID NO:42):

Gene name: prostaglandin G/H synthase 2 (COX-2; PGHS-2)

Unigene number: Hs.196384

Probeset Accession #: D28235

Nucleic Acid Accession #: NM_000963

Coding sequence: 135-1949 (predicted start/stop codons underlined)

Paragraph beginning at line 24 of page 117 has been amended as follows:

ACJ6 DNA sequence (SEQ ID NO:43):

Gene name: SEC14-like-1

Unigene number: Hs.75232

Probeset Accession #: D67029

Nucleic Acid Accession #: NM_003003

Coding sequence: 304-2451 (predicted start/stop codons underlined)

Paragraph beginning at line 56 of page 118 has been amended as follows:

ACJ8 DNA sequence (SEQ ID NO:44):

Gene name: intercellular adhesion molecule 1 (ICAM1; CD54)

Unigene number: Hs.168383

Probeset Accession #: M24283

Nucleic Acid Accession #: NM_000201

Coding sequence: 58-1656 (predicted start/stop codons underlined)

10021650-120501

Paragraph beginning at line 47 of page 119 has been amended as follows:

ACK3 DNA sequence (SEQ ID NO:45):

Gene name: angiopoietin 1 receptor (TIE-2; TEK)

Unigene number: Hs.89640

Probeset Accession #: L06139

Nucleic Acid Accession #: NM_000459

Coding sequence: 149-3523 (predicted start/stop codons underlined)

Paragraph beginning at line 57 of page 120 has been amended as follows:

PZA6 DNA sequence (SEQ ID NO:46):

Gene name: prostate differentiation factor (PLAB; MIC-1)

Unigene number: Hs.116577

Probeset Accession #: AB000584

Nucleic Acid Accession #: NM_004864

Coding sequence: 26-952 (predicted start/stop codons underlined)

Paragraph beginning at line 19 of page 121 has been amended as follows:

AAC8 DNA sequence (SEQ ID NO:47):

Gene name: none

Unigene number: Hs.6682

Probeset Accession #: AA227926

Nucleic Acid Accession #: none

Coding sequence: no ORF identified, possible frameshifts

Paragraph beginning at line 12 of page 122 has been amended as follows:

AAD2 DNA sequence (SEQ ID NO:48):

Gene name: Thrombospondin-1

Unigene number: Hs.87409

Probeset Accession #: AA232645

Nucleic Acid Accession #: NM_003246

Coding sequence: 112-3624 (predicted start/stop codons underlined)

10021660.120601

Paragraph beginning at line 49 of page 123 has been amended as follows:

AAD9 DNA sequence (SEQ ID NO:49):

Gene name: LIM homeobox protein cofactor (CLIM-1)
Unigene number: Hs.4980
Probeset Accession #: F13782
Nucleic Acid Accession #: AF047337
Coding sequence: 110-1231(predicted start/stop codons underlined)

Paragraph beginning at line 29 of page 124 has been amended as follows:

AAE1 DNA sequence (SEQ ID NO:50):

Gene name: guanine nucleotide binding protein 11
Unigene number: Hs.83381
Probeset Accession #: U31384
Nucleic Acid Accession #: NM_004126.1
Coding sequence: 108-329 (predicted start/stop codons underlined)

Paragraph beginning at line 49 of page 124 has been amended as follows:

AAE2 DNA sequence (SEQ ID NO:51):

Gene name: Transcription factor 4 (immunoglobulin transcription factor 2) (ITF-2) (SL3-3 Enhancer factor 2) (SEF-2)
Unigene number: Hs.289068
Probeset Accession #: M74719
Nucleic Acid Accession #: NM_003199.1
Coding sequence: 200-2203 (predicted start/stop codons underlined)

Paragraph beginning at line 33 of page 125 has been amended as follows:

AAE4 DNA sequence (SEQ ID NO:52):

Gene name: phosphatidylcholine 2-acylhydrolase
Unigene number: Hs.211587
Probeset Accession #: M68874
Nucleic Acid Accession #: M68874
Coding sequence: 139-2388 (predicted start/stop codons underlined)

10021560-120601

Paragraph beginning at line 22 of page 126 has been amended as follows:

ACA1 DNA sequence (SEQ ID NO:53):

Gene name: tissue factor pathway inhibitor 2 TFPI2, placental protein 5 (PP5)
Unigene number: Hs.78045
Probeset Accession #: D29992
Nucleic Acid Accession #: D29992.1
Coding sequence: 57-764 (predicted start/stop codons underlined)

Paragraph beginning at line 51 of page 126 has been amended as follows:

ACB8 DNA sequence (SEQ ID NO:54):

Gene name: myosin X
Unigene number: Hs.61638
Probeset Accession #: N77151
Nucleic Acid Accession #: NM_012334
Coding sequence: 223-6399 (predicted start/stop codons underlined)

Paragraph beginning at line 54 of page 128 has been amended as follows:

ACC3 DNA sequence (SEQ ID NO:55):

Gene name: calcitonin receptor-like (CALCRL)
Unigene number: Hs.152175
Probeset Accession #: L76380
Nucleic Acid Accession #: NM_005795
Coding sequence: 555-1940 (predicted start/stop codons underlined)

Paragraph beginning at line 46 of page 129 has been amended as follows:

ACC4 DNA sequence (SEQ ID NO:56):

Gene name: Homo sapiens mRNA; cDNA DKFZp586E1624
Unigene number: Hs.94030
Probeset Accession #: AA452000
Nucleic Acid Accession #: AL110152.1
Coding sequence: no ORF identified, possible frameshifts

10021660-120601

Paragraph beginning at line 10 of page 130 has been amended as follows:

ACC5 DNA sequence (SEQ ID NO:57):

Gene name: Selectin E (endothelial adhesion molecule 1)

Unigene number: Hs.89546

Probeset Accession #: M24736

Nucleic Acid Accession #: NM_000450

Coding sequence: 117-1949 (predicted start/stop codons underlined)

Paragraph beginning at line 15 of page 131 has been amended as follows:

ACC8 DNA sequence (SEQ ID NO:58):

Gene name: Chemokine (C-X-C motif), receptor 4 (fusin)

Unigene number: Hs.89414

Probeset Accession #: L06797

Nucleic Acid Accession #: NM_003467

Coding sequence: 89-1147 (predicted start/stop codons underlined)

Paragraph beginning at line 52 of page 131 has been amended as follows:

ACF2 DNA sequence (SEQ ID NO:59):

Gene name: Endothelial cell-specific molecule 1

Unigene number: Hs.41716

Probeset Accession #: X89426

Nucleic Acid Accession #: NM_007036

Coding sequence: 56-610 (predicted start/stop codons underlined)

Paragraph beginning at line 27 of page 132 has been amended as follows:

ACF4 DNA sequence (SEQ ID NO:60):

Gene name: P53-responsive gene 2 similar to D.melanogaster peroxidasin(U11052)

Unigene number: Hs.118893

Probeset Accession #: D86983

Nucleic Acid Accession #: D86983

Coding sequence: 1-4491 (predicted stop codon underlined, sequence is open at 5' end)

10021660.120601

Paragraph beginning at line 61 of page 133 has been amended as follows:

ACF5 DNA sequence (SEQ ID NO:61):

Gene name: Mitogen-activated protein kinase kinase kinase kinase 4
Unigene number: Hs.3628
Probeset Accession #: N54067
Nucleic Acid Accession #: NM_004834
Coding sequence: 80-3577 (predicted start/stop codons underlined)

Paragraph beginning at line 67 of page 134 has been amended as follows:

ACF8 DNA sequence (SEQ ID NO:62):

Gene name: Phospholipase A2, group IVC (cytosolic, calcium-independent)
Unigene number: Hs.18858
Probeset Accession #: AA054087
Nucleic Acid Accession #: NM_003706
Coding sequence: 310-1935 (predicted start/stop codons underlined)

Paragraph beginning at line 50 of page 135 has been amended as follows:

ACG1 DNA sequence (SEQ ID NO:63):

Gene name: Carbohydrate (chondroitin 6/keratan) sulfotransferase 1
Unigene number: Hs.104576
Probeset Accession #: AA868063
Nucleic Acid Accession #: NM_003654
Coding sequence: 367-1602 (predicted start/stop codons underlined)

Paragraph beginning at line 32 of page 136 has been amended as follows:

ACG5 DNA sequence (SEQ ID NO:64):

Gene name: Multimerin
Unigene number: Hs.268107
Probeset Accession #: U27109
Nucleic Acid Accession #: U27109.1
Coding sequence: 72-3758 (predicted start/stop codons underlined)

10021650.120601

Paragraph beginning at line 43 of page 137 has been amended as follows:

ACC6 DNA sequence (SEQ ID NO:65):

Gene name: Homo sapiens cDNA FLJ11502 fis, clone HEMBA1002102, weakly similar to ANKRYIN

Unigene number: Hs.213194

Probeset Accession #: AA187101

Nucleic Acid Accession #: AK021564

Coding sequence: 1-450 (predicted stop codon underlined, 5'end sequence is open)

Paragraph beginning at line 13 of page 138 has been amended as follows:

ACC7 DNA sequence (SEQ ID NO:66):

Gene name: Human RAL A gene

Unigene number: Hs.6906

Probeset Accession #: AA083572

Nucleic Acid Accession #: contig of X15014.1 and AK026850

Coding sequence: 1-621 (predicted start/stop codons underlined)

Paragraph beginning at line 65 of page 138 has been amended as follows:

ACC9 DNA sequence (SEQ ID NO:67):

Gene name: KIAA0955 protein

Unigene number: Hs.10031

Probeset Accession #: AA027168

Nucleic Acid Accession #: AB023172

Coding sequence: 314-1609 (predicted start/stop codons underlined)

Paragraph beginning at line 23 of page 140 has been amended as follows:

ACF6 DNA sequence (SEQ ID NO:68):

Gene name: Homo sapiens cDNA FLJ10669 fis, clone NT2RP2006275, weakly similar to Microtubule-associated protein 1B [CONTAINS: LIGHT CHAIN LC1]

Unigene number: Hs.66048

Probeset Accession #: AA609717

Nucleic Acid Accession #: AK001531

Coding sequence: 176-2194 (predicted start/stop codons underlined)

10021560.120504

Paragraph beginning at line 4 of page 141 has been amended as follows:

AAA4 Protein sequence+ (SEQ ID NO:69):

Gene name: CGI-100 protein

Unigene number: Hs.275253

Probeset Accession #: AA089688

Protein Accession #: NP_057124

Signal sequence: predicted 1-23 (first underlined sequence)

Transmembrane Domain: predicted 201-217 (second underlined sequence)

emp24/gp25L/p24 domain: predicted 13-227

Summary: gp25L/emp24/p24 protein family members of the cis-Golgi network bind both COP I and II coatomer. Members of this family are implicated in bringing cargo forward from the ER and binding to coat proteins by their cytoplasmic domains.

Paragraph beginning at line 22 of page 141 has been amended as follows:

AAA7 Protein sequence+ (SEQ ID NO:70):

Gene name: Endothelial differentiation, sphingolipid G-protein-coupled receptor, 1 (EDG1)

Unigene number: Hs.154210

Probeset Accession #: M31210

Protein Accession#: NP_001391

7 Transmembrane Domains: predicted 50-71, 92-110, 122-140, 160-177, 201-222, 251-269, 281-301 (underlined sequences)

Summary: Endothelial differentiation, sphingolipid G-protein-coupled receptor, 1 may regulate the differentiation of endothelial cells. It binds the sphingolipid metabolite, sphingosine-1-phosphate, which may function as a second messenger in cell proliferation and survival.

Paragraph beginning at line 44 of page 141 has been amended as follows:

AAB3 Protein sequence+ (SEQ ID NO:71)

Gene name: Solute carrier family 20 (phosphate transporter), member 1, Human leukaemia virus receptor 1 (GLVR1)

Unigene number: Hs.78452

Probeset Accession #: L20859

Protein Accession #: NP_005406

Transmembrane domains: predicted 24-40, 62-78, 164-180, 198-214, 232-248, 513-529, 562-578, 604-620, 655-671

Cellular Localization: Likely a Type IIIa membrane protein (Ncyt Cexo)

10021560 120601

Paragraph beginning at line 68 of page 141 has been amended as follows:

AAB4 Protein sequence+ (SEQ ID NO:72):
Gene name: Matrix metalloproteinase 10 (stromelysin 2)
Unigene number: Hs.2258
Probeset Accession #: X07820
Protein Accession #: NP_002416
Signal sequence: predicted 1-17 (underlined sequence)
Cellular Localization: predicted secreted

Paragraph beginning at line 18 of page 142 has been amended as follows:

AAB6 Protein sequence (SEQ ID NO:73):
Gene name: Podocalyxin-like
Unigene number: Hs.16426
Probeset Accession #: U97519
Protein Accession #: NP_005388
Transmembrane domain: predicted 432-448 (underlined sequence)
Cellular Localization: predicted Type Ia membrane protein (Nexo)

Paragraph beginning at line 37 of page 142 has been amended as follows:

AAB8 Protein sequence+ (SEQ ID NO:74):
Gene name: EGF-containing fibulin-like extracellular matrix protein 1
Unigene number: Hs.76224
Probeset Accession #: U03877
Protein Accession #: NP_004096 Variant 1
Signal sequence: predicted 1-17 (underlined sequence)
Summary: This gene spans approximately 18 kb of genomic DNA and consists of 12 exons. Two transcripts with distinct 5' UTR have been described; the resulting proteins have distinct N-terminal amino acid sequences. Translation initiation from internal methionine residues was observed with in vitro translation. A signal peptide sequence is predicted for translation initiation sites 1, 2, and 4. The protein isoforms contain 5 or 6 calcium-binding EGF2 domains and 5 or 6 EGF2 domains. Mutations in this gene cause the retinal disease Malattia Leventinese. Transcript Variant: This variant (1) has a distinct 5' UTR and N-terminal protein sequence as compared to variant 2.

10021560 120601

Paragraph beginning at line 64 of page 142 has been amended as follows:

AAB9 Protein sequence+ (SEQ ID NO:75):

Gene name: Melanoma adhesion molecule, MUC 18 glycoprotein
Unigene number: Hs.211579
Probeset Accession #: M28882
Protein Accession #: NP_006491
Signal sequence: predicted 1-17 (first underlined sequence)
Transmembrane domain: predicted 559-575 (second underlined sequence)
Cellular localization: predicted Type Ia membrane protein (Nexo)

Paragraph beginning at line 18 of page 143 has been amended as follows:

AAC1 Protein sequence+ (SEQ ID NO:76):

Gene name: Matrix metalloproteinase 1 (interstitial collagenase)
Unigene number: Hs.83169
Probeset Accession #: X54925
Protein Accession #: NP_002412
Signal sequence: predicted 1-19 (underlined sequence)
Cellular localization: predicted secreted protein

Paragraph beginning at line 36 of page 143 has been amended as follows:

AAC3 Protein sequence+ (SEQ ID NO:77):

Gene name: Branched chain aminotransferase 1, cytosolic
Unigene number: Hs.157205
Probeset Accession #: AA423987
Protein Accession #: NP_005495
Cellular Localization: cytoplasmic
Summary: The lack of the cytosolic enzyme branched-chain amino acid transaminase (BCT) causes cell growth inhibition. There may be at least 2 different clinical disorders due to a defect of branched-chain amino acid transamination: hypervalinemia and hyperleucine-isoleucinemia. Since there are 2 distinct BCATs, mitochondrial and cytosolic, it is possible that one is mutant in each of these 2 conditions.

10021660.12060.1

Paragraph beginning at line 58 of page 143 has been amended as follows:

ACG4 Protein sequence+ (SEQ ID NO:78):

Gene name: Pentaxin-related gene, rapidly induced by IL-1 beta

Unigene number: Hs.2050

Probeset Accession #: M31166

Protein Accession #: NP_002843

Signal sequence: predicted 1-17 (underlined sequence)

Cellular localization: predicted secreted

Summary: TNF-inducible member of hyaluronate binding protein family, related to CD44

Paragraph beginning at line 9 of page 144 has been amended as follows:

ACK5 Protein sequence+ (SEQ ID NO:79):

Gene name: Von Willebrand factor; Coagulation factor VIII

Unigene number: Hs.110802

Probeset Accession #: M10321

Protein Accession #: NP_000543

Signal peptide: predicted 1-22 (underlined sequence)

Cellular localization: predicted secreted

Paragraph beginning at line 66 of page 144 has been amended as follows:

AAC7 protein sequence+ (SEQ ID NO:80):

Gene name: KIAA1294 protein

Probeset Accession #: AA432248

Protein Accession #: BAA92532

Cellular localization: predicted nuclear protein

PFAM prediction: 22-153 Band 41 domain (underlined seq). A number of cytoskeletal-associated proteins that associate with various proteins at the interface between the plasma membrane and the cytoskeleton contain a conserved N-terminal domain of about 150 amino-acid residues.

Paragraph beginning at line 28 of page 145 has been amended as follows:

ACG8 Protein sequence+ (SEQ ID NO:81):

Gene name: ubiquitin E3 ligase SMURF2

Unigene number: Hs.21806 (3'UTR only)

Probeset Accession #: AA398243

Protein Accession #: AF301463_1

Cellular Localization: predicted cytoplasmic

Summary: Smurf2 Is a Ubiquitin E3 Ligase Mediating Proteasome-dependent Degradation of Smad2 in Transforming Growth Factor-beta Signaling

10021660-120601

Paragraph beginning at line 52 of page 145 has been amended as follows:

ACH1 Protein sequence+ (SEQ ID NO:82):

Gene name: EST
Unigene number: Hs.30089
Probeset Accession #: AA410480
CAT cluster#: cluster 96816_1
Summary: predicted open reading frame

Paragraph beginning at line 65 of page 145 has been amended as follows:

ACJ2 Protein sequence+ (SEQ ID NO:83):

Gene name: Complement component Clq receptor
Unigene number: Hs.97199
Probeset Accession #: AA487558
Protein Accession #: NP_036204
Signal sequence: 1-17 (first underlined sequence)
Transmembrane domain: 589-605 (second underlined sequence)
Cellular localization: This gene encodes a predicted type I membrane protein.
Summary: This protein acts as a receptor for complement protein Clq, mannose-binding lectin, and pulmonary surfactant protein A. This protein is a functional receptor involved in ligand-mediated enhancement of phagocytosis.

Paragraph beginning at line 22 of page 146 has been amended as follows:

ACJ3 Protein sequence+ (SEQ ID NO:84):

Gene name: FLT1/vascular endothelial growth factor receptor
Unigene number: Hs.138671
Probeset Accession #: AA047437
Transmembrane domain: predicted 764-780 (underlined sequence)
Cellular Localization: predicted cell surface tyrosine kinase

Paragraph beginning at line 54 of page 146 has been amended as follows:

ACJ9 Protein sequence+ (SEQ ID NO:85):

Gene name: Purine nucleoside phosphorylase
Unigene number: Hs.75514
Probeset Accession #: K02574
Protein Accession #: CAA25320
Cellular Localization: predicted cytoplasmic
Summary: likely to catalyze the reversible phosphorolytic cleavage of purine ribonucleosides and 2'-deoxyribonucleosides

10021560.120601

Paragraph beginning at line 2 of page 147 has been amended as follows:

ACK4 Protein sequence (SEQ ID NO:86):

Gene name: EST

Probeset Accession #: R68763

Predicted amino acid seq: FGENESH exon prediction on BAC clone AC009414

Predicted nuclear target motifs: from 25 (4) RRRP (SEQ ID NO:87) (underlined);

176 (5) RRRR (SEQ ID NO:88) (underlined); 177 (5) RRRR (SEQ ID NO:88)

(underlined); 239 (5) KRKK (SEQ ID NO:89) (underlined); 399 (4) PPRARRT (SEQ ID

NO:90) (underlined); 400 (5) PRARRTE (SEQ ID NO:91) (underlined)

Cellular localization: predicted nuclear

Paragraph beginning at line 22 of page 147 has been amended as follows:

AAA8 Protein sequence (SEQ ID NO:92):

Gene name: ETL protein, with extended open reading frame

Unigene number: Hs.57958

Probeset Accession #: D58024

Protein Accession #: AAG33021

Transmembrane domains: predicted 454-470, 486-502, 511-527, 528-544, 556-572, 600-616, 642-661, 672-689 (underlined sequences)

Extended sequence: Residues 1-564 were added to the sequence in AAG33021

Cellular Localization: predicted cell surface serpentine receptor

Paragraph beginning at line 46 of page 147 has been amended as follows:

AAC6 Protein sequence+ (SEQ ID NO:93):

Gene name: EST

Unigene number: Hs.134797

Probeset Accession #: AA025351

Protein accession #: BAB14599

Signal sequence: predicted 1-24 (first underlined sequence)

extended sequence: second underlined sequence

Paragraph beginning at line 4 of page 148 has been amended as follows:

ACH7 Protein sequence+ (SEQ ID NO:94):

Gene name: EST

Unigene number: Hs.3807

Probeset Accession #: AA292694

BAC Accession #: AL161751

FGENESH predicted aa seq: 1-647; based on BAC clone AL161751

10021660.120601

Paragraph beginning at line 24 of page 148 has been amended as follows:

AAD4 Protein sequence (SEQ ID NO:95):

Gene name: ERG
Unigene number: Hs.45514
Probeset Accession #: R32894
Protein Accession #: AAA52398
Signal sequence: none
Transmembrane domains: none
PFAM domains: predicted Ets-domain 294-373; SAM_PNT: 122-206
Summary: ERG2 is a sequence-specific DNA-binding protein.

Paragraph beginning at line 44 of page 148 has been amended as follows:

AAD5 Protein sequence (SEQ ID NO:96):

Gene name: activin A receptor type II-like 1 (ALK-1)
Unigene number: Hs.172670
Probeset Accession #: T57112
Protein Accession #: NP_000011
Signal sequence: predicted 1-21
Transmembrane domain: predicted 119-135
PFAM domains: predicted pkinase 204-489
Summary: Type Ia membrane protein; receptor tyrosine kinase

Paragraph beginning at line 21 of page 149 has been amended as follows:

ACA4 Protein sequence (SEQ ID NO:97):

Gene name: alpha satellite junction DNA sequence
Unigene number: Hs.247946
Probeset Accession #: M21305
Protein Accession #: AAA88020
Signal sequence: none
Transmembrane domains: none
PFAM domains: none

TD9021"09512001

Paragraph beginning at line 33 of page 149 has been amended as follows:

ACG6 Protein sequence (SEQ ID NO:98):

Gene name: intercellular adhesion molecule 2 (ICAM2)

Unigene number: Hs.83733

Probeset Accession #: M32334

Protein Accession #: NP_000864

Signal sequence: predicted 1-21

Transmembrane domain: predicted 224-248

PFAM domains: predicted 41-98, 127-197; immunoglobulin-like C2-type domains

Summary: a predicted Type Ia membrane protein; it plays a role in cell adhesion and is the ligand for the LFA-1 protein. ICAM2 is also called CD102.

Paragraph beginning at line 51 of page 149 has been amended as follows:

ACG7 Protein sequence (SEQ ID NO:99):

Gene name: Cadherin 5, VE-cadherin (CDH5)

Unigene number: Hs.76206

Probeset Accession #: X79981

Protein Accession #: NP_001786

Signal sequence: predicted 1-27

Transmembrane domain: predicted 604-620

PFAM domains: Cadherin domains predicted 53-141, 156-249, 263-364, 377-470, and 487-576

Summary: Likely a Type I membrane protein. Cadherins are calcium-dependent adhesive proteins that mediate cell-to-cell interaction. VE-cadherin is associated with intercellular junctions.

Paragraph beginning at line 12 of page 150 has been amended as follows:

ACG9 Protein sequence (SEQ ID NO:100):

Gene name: lysyl oxidase-like 2 (LOXL2)

Unigene number: Hs.83354

Probeset Accession #: U89942

Protein Accession #: NP_002309

Signal sequence: predicted 1-25

Transmembrane domains: none predicted

PFAM domains: scavenger receptor cysteine-rich domains predicted 68-159, 203-238, 336-425, 439-528; Lysyl oxidase predicted 548-749.

Summary: Likely a secreted protein. Lysyl oxidase is a copper-dependent amine oxidase that belongs to a heterogeneous family of enzymes that oxidize primary amine substrates to reactive aldehydes, acting on the extracellular matrix substrates, e.g., collagen and elastin.

10021660 120601
T0902T 099T200T

Paragraph beginning at line 41 of page 150 has been amended as follows:

ACH2 Protein sequence (SEQ ID NO:101):

Gene name: TIE tyrosine-protein kinase
Unigene number: Hs.78824
Probeset Accession #: X60957
Protein Accession #: NP_005415
Signal sequence: predicted 1-21
Transmembrane domain: predicted 770-786
PFAM domains: laminin-EGF predicted 234-267; FN3 predicted 460-520, 548-632, and 644-729; tyrosine_kinase predicted 839-1107
Summary: Likely a Type Ia membrane protein; TIE is a tyrosine-kinase receptor with an unknown ligand; its expression is likely necessary for normal blood vessel development.

Paragraph beginning at line 7 of page 151 has been amended as follows:

ACH3 Protein sequence (SEQ ID NO:102):

Gene name: placental growth factor (PGF; PlGF1; VEGF-related protein)
Unigene number: Hs.2894
Probeset Accession #: X54936
Protein Accession #: NP_002623
Signal sequence: predicted 1-21
Transmembrane domain: none predicted
PFAM domains: PDGF predicted 52-130
Summary: Likely a secreted protein; likely regulates angiogenesis by interacting with FLT1 and FLK1.

Paragraph beginning at line 23 of page 151 has been amended as follows:

ACH4 Protein sequence (SEQ ID NO:103):

Gene name: nidogen 2 (NID2)
Unigene number: Hs.82733
Probeset Accession #: D86425
Protein Accession #: NP_031387
Signal sequence: predicted 1-30
Transmembrane domain: none predicted
PFAM domains: EGF-like_domains predicted 489-524, 764-800, 806-843, 853-891, and 897-930; thyroglobulin_repeats predicted 941-1006, and 1020-1085; LDL_receptor_repeats predicted 1155-1197, 1199-1240, and 1242-1285.
Summary: A secreted protein; NID2 likely interacts with collagens I and IV and laminin-1 to promote cell adhesion to the basement membrane.

10021660-120601

Paragraph beginning at line 61 of page 151 has been amended as follows:

ACH5 Protein sequence (SEQ ID NO:104):

Gene name: SNL (singled-like; sea urchin fascin homolog-like)

Unigene number: Hs.118400

Probeset Accession #: U03057

Protein Accession #: NP_003079

Signal sequence: none identified

Transmembrane domain: none identified

PFAM domains: none identified

Summary: a cytoplasmic, actin-bundling protein that is likely to be involved in the assembly of actin filament bundles present in microspikes, membrane ruffles, and stress fibers

Paragraph beginning at line 16 of page 152 has been amended as follows:

ACH6 Protein sequence (SEQ ID NO:105):

Gene name: endothelial protein C receptor (EPCR; PROCR)

Unigene number: Hs.82353

Probeset Accession #: L35545

Protein Accession #: NP_006395

Signal sequence: predicted 1-17

Transmembrane domain: predicted 211-227

PFAM domains: none identified

Summary: a Type Ia membrane protein, EPCR likely binds to [thrombin]-activated Protein C, a vitamin K-dependent serine protease zymogen necessary for blood coagulation.

Paragraph beginning at line 34 of page 152 has been amended as follows:

ACH8 Protein sequence (SEQ ID NO:106):

Gene name: melanoma adhesion molecule (MCAM; MUC18)

Unigene number: Hs.211579

Probeset Accession #: D51069

Protein Accession #: NP_006491

Signal sequence: predicted 1-17

Transmembrane domain: predicted 559-575

PFAM domains: immunoglobulin_domains predicted 264-324, and 356-410.

Summary: a Type Ia membrane protein, associated with tumor progression and the development of metastasis in human malignant melanoma, and may play a role in neural crest cells during embryonic development.

10021560.120601

Paragraph beginning at line 59 of page 152 has been amended as follows:

ACH9 Protein sequence (SEQ ID NO:107):

Gene name: endothelin-1 (EDN1)

Unigene number: Hs.2271

Probeset Accession #: J05008

Protein Accession #: NP_001946

Signal sequence: predicted 1-17

Transmembrane domain: none predicted

PFAM domains: Endothelin domains predicted 59-73, and 108-129.

Summary: a secreted zymogen; the active protein is likely a 26-amino acid peptide with potent mammalian vasoconstrictor activity; it is necessary for normal vessel development.

Paragraph beginning at line 11 of page 153 has been amended as follows:

ACJ1 Protein sequence (SEQ ID NO:108):

Gene name: BMX non-receptor tyrosine kinase

Unigene number: Hs.27372

Probeset Accession #: X83107

Protein Accession #: NP_001712

Signal sequence: none identified

Transmembrane domain: none identified

PFAM domains: plektrin_homology_domain predicted 6-111; SH2_domain predicted 294-383; protein_kinase_domain predicted 417-663

Summary: a cytoplasmic protein, it likely plays a role in the growth and differentiation of hematopoietic cells; it is known to also be expressed in endothelial cells.

Paragraph beginning at line 38 of page 153 has been amended as follows:

ACJ4 Protein sequence (SEQ ID NO:109):

Gene name: prostaglandin G/H synthase 2 (COX-2; PGHS-2)

Unigene number: Hs.196384

Probeset Accession #: D28235

Protein Accession #: NP_000954

Signal sequence: predicted 1-17

Transmembrane domain: none identified

PFAM domains: EGF-like_domain predicted 18-55.

Summary: a microsomal enzyme; COX-2 is the therapeutic target of the nonsteroidal anti-inflammatory drugs (NSAIDs), such as aspirin.

10021660.120601

Paragraph beginning at line 62 of page 153 has been amended as follows:

ACJ6 Protein sequence (SEQ ID NO:110):

Gene name: SEC14-like-1
Unigene number: Hs.75232
Probeset Accession #: D67029
Protein Accession #: NP_002994
Signal sequence: none identified
Transmembrane domain: none identified
PFAM domains: none identified
Summary: a cytoplasmic protein

Paragraph beginning at line 18 of page 154 has been amended as follows:

ACJ8 Protein sequence (SEQ ID NO:111):

Gene name: intercellular adhesion molecule 1 (ICAM1; CD54)
Unigene number: Hs.168383
Probeset Accession #: M24283
Protein Accession #: NP_000192
Signal sequence: predicted 1-27
Transmembrane domain: predicted 481-497
PFAM domains: immunoglobulin_domains predicted 128-188, and 325-373.
Summary: a Type 1a membrane protein; ICAM1 is typically expressed on endothelial cells and cells of the immune system; ICAM1 binds to integrins of type CD11a/CD18, or CD11b/CD18; ICAM1 is also exploited by Rhinovirus as a receptor.

Paragraph beginning at line 41 of page 154 has been amended as follows:

ACK3 Protein sequence (SEQ ID NO:112):

Gene name: angiopoietin 1 receptor (TIE-2; TEK)
Unigene number: Hs.89640
Probeset Accession #: L06139
Protein Accession #: NP_000450
Signal sequence: predicted 1-18
Transmembrane domain: predicted 746-770
PFAM domains: immunoglobulin_domains predicted 44-102, 370-424; EGF_like_domains predicted 210-252, 254-299, and 301-341; FN3_domains predicted 444-536, 541-634, and 638-732; protein_kinase_domain predicted 824-1096.
Summary: a Type 1a membrane protein; it is expressed almost exclusively in endothelial cells in mice, rats, and humans; the ligand for this receptor is angiopoietin-1; defects in TEK are associated with inherited venous malformations; the TEK signaling pathway appears to be critical for endothelial cell-smooth muscle cell communication in venous morphogenesis.

10021650-120601
FOIA b 7 - D

Paragraph beginning at line 10 of page 155 has been amended as follows:

PZA6 Protein sequence (SEQ ID NO:113):

Gene name: prostate differentiation factor (PLAB; MIC-1)

Unigene number: Hs.116577

Probeset Accession #: AB000584

Protein Accession #: NP_004855

Signal sequence: predicted 1-29

Transmembrane domain: none identified

PFAM domains: TGF-beta _domain predicted 211-308.

Summary: a secreted protein; its exact function is unclear; it inhibits proliferation of primitive hematopoietic progenitors; it inhibits activation of macrophages; it is highly expressed in placenta and in serum of pregnant women; it may promote fetal survival by suppressing the production of maternally-derived proinflammatory cytokines within the uterus.

Paragraph beginning at line 32 of page 155 has been amended as follows:

AAD2 Protein sequence+ (SEQ ID NO:114):

Gene name: Thrombospondin-1

Unigene number: Hs.87409

Probeset Accession #: AA232645

Protein Accession #: NP_003237.1

Signal sequence: predicted 1-18 (first underlined sequence)

Transmembrane Domain: none identified

Summary: Thrombospondin is a large modular glycoprotein component of the extracellular matrix and contains a variety of distinct domains, including three repeating subunits (types I, II, and III) that share homology to an assortment of other proteins.

Paragraph beginning at line 66 of page 155 has been amended as follows:

AAD9 protein sequence (SEQ ID NO:115):

Gene name: LIM homeobox protein cofactor (CLIM-1)

Unigene number: Hs.4980

Probeset Accession #: F13782

Protein Accession #: AAC83552

Pfam: LIM bind

Transmembrane Domain: none identified

Summary: The LIM homeodomain (LIM-HD) proteins, which contain two tandem LIM domains followed by a homeodomain, are critical transcriptional regulators of embryonic development. The LIM domain is a conserved cysteine-rich zinc-binding motif found in LIM-HD proteins, cytoskeletal components, LIM kinases, and other proteins. LIM domains are protein-protein interaction motifs, can inhibit binding of LIM-HD proteins to DNA, and can negatively regulate LIM-HD protein function.

10021650-120501

Paragraph beginning at line 21 of page 156 has been amended as follows:

AAE1 protein sequence (SEQ ID NO:116):

Gene name: guanine nucleotide binding protein 11

Unigene number: Hs.83381

Probeset Accession #: U31384

Protein Accession #: NP_004117.1

Pfam: G-gamma; CAAX motif (farnesylation site) prediction underlined

Summary: The G gamma proteins are a component of the trimeric G-proteins that interact with cell surface receptors. The G protein beta and gamma subunits directly regulate the activities of various enzymes and ion channels after receptor ligation. Unlike most of the other known gamma subunits, gamma 11 is modified by a farnesyl group and is not capable of interacting with beta 2.

Paragraph beginning at line 37 of page 156 has been amended as follows:

AAE2 protein sequence (SEQ ID NO:117):

Gene name: Transcription factor 4 (Immunoglobulin transcription factor 2) (ITF-2) (SL3-3 Enhancer factor 2) (SEF-2)

Unigene number: Hs.289068

Probeset Accession #: M74719

Protein Accession #: NP_003190.1

Pfam: HLH domain prediction underlined

Summary: Transcription factor 4 is a helix-loop-helix (HLH) protein which belongs to a family of nuclear proteins, designated SL3-3 enhancer factors 2 (SEF2), that interact with an Ephrussi box-like motif within the glucocorticoid response element in the enhancer of the murine leukemia virus SL3-3. Various cell types display differences both in the sets of SEF2-DNA complexes formed and in their amounts. Molecular analysis of cDNA clones show the existence of multiple related mRNA species containing alternative coding regions, which are most probably a result of differential splicing.

Paragraph beginning at line 68 of page 156 has been amended as follows:

AAE4 protein sequence (SEQ ID NO:118):

Gene name: phosphatidylcholine 2-acylhydrolase

Unigene number: Hs.211587

Probeset Accession #: M68874

Protein Accession #: AAA60105.1

Pfam: PLA2 B, C2 domain prediction underlined

Summary: Phospholipases A2 (PLA2s) play a key role in inflammatory processes through production of precursors of eicosanoids and platelet-activating factor. PLA2 is a 100 kd protein that contains a structural element homologous to the C2 region of protein kinase C.

10021660-120601

Paragraph beginning at line 26 of page 157 has been amended as follows:

ACA1 protein sequence (SEQ ID NO:119):

Gene name: tissue factor pathway inhibitor 2 TFPI2, placental protein 5 (PP5)

Unigene number: Hs.78045

Probeset Accession #: D29992

Protein Accession #: BAA06272.1

Pfam: Kunitz BPTI

Signal sequence: underlined

Summary: ACA1 is a serine proteinase inhibitor that was originally purified from conditioned medium of the human glioblastoma cell line T98G. ACA1 is identical to placental protein 5 (PP5) and TFPI2, a placenta-derived glycoprotein with serine proteinase inhibitor activity. PP5 belongs to the Kunitz-type serine proteinase inhibitor family, having three putative Kunitz-type inhibitor domains.

Paragraph beginning at line 45 of page 157 has been amended as follows:

ACB8 protein sequence (SEQ ID NO:120):

Gene name: myosin X

Unigene number: Hs.61638

Probeset Accession #: N77151

Protein Accession #: NP_036466

Pfam: myosin head, IQ (calmodulin binding motif), PH, MyTH4

Summary: Myosins are molecular motors that move along filamentous actin. Seven classes of myosin are expressed in vertebrates: conventional myosin, or myosin-II, as well as the 6 unconventional myosin classes-I, -V, -VI, -VII, -IX, and -X.

Paragraph beginning at line 24 of page 158 has been amended as follows:

ACC3 protein sequence (SEQ ID NO:121):

Gene name: calcitonin receptor-like (CALCRL)

Unigene number: Hs.152175

Probeset Accession #: L76380

Protein Accession #: NP_005786.1

Pfam: 7TM 2 (7 transmembrane receptor (Secretin family))

Transmembrane domains: predictions underlined

Signal sequence: first underlined region

Summary: Calcitonin gene-related peptide (CGRP) is a neuropeptide with diverse biological effects including potent vasodilator activity. The human CGRP1 receptor shares significant peptide sequence homology with the human calcitonin receptor, a member of the G-protein-coupled receptor superfamily. Stable expression in 293 (HEK 293) cells produces specific, high affinity binding sites for CGRP. Exposure of these cells to CGRP results in a 60-fold increase in cAMP production.

10021650-120601

Paragraph beginning at line 49 of page 158 has been amended as follows:

ACC5 protein sequence (SEQ ID NO:122):

Gene name: Selectin E (endothelial adhesion molecule 1)

Unigene number: Hs.89546

Probeset Accession #: M24736

Protein Accession #: NP_000441.1

Pfam: lectin c, EGF like domain, sushi (SCR domain)

Signal sequence: first underlined region

Transmembrane domain: second underlined region

Summary: Focal adhesion of leukocytes to the blood vessel lining is a key step in inflammation and certain vascular disease processes. Endothelial leukocyte adhesion molecule-1 (ELAM-1), a cell surface glycoprotein expressed by cytokine-activated endothelium, mediates the adhesion of blood neutrophils. The primary sequence of ELAM-1 predicts an amino-terminal lectin-like domain, an EGF domain, and six tandem repetitive motifs (about 60 amino acids each) related to those found in complement regulatory proteins. A similar domain structure is also found in the MEL-14 lymphocyte cell surface homing receptor, and in granule-membrane protein 140, a membrane glycoprotein of platelet and endothelial secretory granules that can be rapidly mobilized (less than 5 minutes) to the cell surface by thrombin and other stimuli. Thus, ELAM-1 may be a member of a nascent gene family of cell surface molecules involved in the regulation of inflammatory and immunological events at the interface of vessel wall and blood.

Paragraph beginning at line 17 of page 159 has been amended as follows:

ACC8 protein sequence (SEQ ID NO:123):

Gene name: Chemokine (C-X-C motif), receptor 4 (fusin)

Unigene number: Hs.89414

Probeset Accession #: L06797

Protein Accession #: NP_003458.1

Pfam: 7TM 1 (7 transmembrane receptor (rhodopsin family))

Signal sequence: none identified

Transmembrane domains: predictions underlined

Summary: The chemokine receptor CXCR4 (also designated fusin and LESTR) is a cofactor for fusion and entry of T cell-tropic strains of HIV-1.

10021660.12060.1

Paragraph beginning at line 36 of page 159 has been amended as follows:

ACF2 protein sequence (SEQ ID NO:124):

Gene name: Endothelial cell-specific molecule 1

Unigene number: Hs.41716

Probeset Accession #: X89426

Protein Accession #: NP_008967.1

Signal sequence: underlined

Pfam: IGFBP (Insulin-like growth factor binding proteins)

Summary: Human endothelial cell-specific molecule (called ESM-1) was cloned from a human umbilical vein endothelial cell (HUVEC) cDNA library. Constitutive ESM-1 gene expression is seen in HUVECs but not in the other human cell lines. The cDNA sequence contains an open reading frame of 552 nucleotides and a 1398-nucleotide 3'-untranslated region including several domains involved in mRNA instability and five putative polyadenylation consensus sequences. The deduced 184-amino acid sequence defines a cysteine-rich protein with a functional NH2-terminal hydrophobic signal sequence.

Paragraph beginning at line 58 of page 159 has been amended as follows:

ACF4 protein sequence (SEQ ID NO:125):

Gene name: P53-responsive gene 2 similar to D.melanogaster peroxidase(U11052)

Unigene number: Hs.118893

Probeset Accession #: D86983

Protein Accession #: BAA13219

Pfam: LRRNT (Leucine rich repeat N-terminal domain), LRR (Leucine Rich Repeat), LRRCT (Leucine rich repeat C-terminal domain), Ig (immunoglobulin domain), Peroxidase, VWC (von Willebrand factor type C domain)

Summary: ACF4 is a gene originally identified from KG-1 cell and brain cDNA libraries.

10021660-120601

Paragraph beginning at line 28 of page 160 has been amended as follows:

ACF5 protein sequence (SEQ ID NO:126):

Gene name: Mitogen-activated protein kinase kinase kinase kinase 4

Unigene number: Hs.3628

Probeset Accession #: N54067

Protein Accession #: NP_004825.1

Pfam: pkinase (Eukaryotic protein kinase domain), CNH domain

Summary: The yeast serine/threonine kinase STE20 activates a signaling cascade that includes STE11 (mitogen-activated protein kinase kinase kinase), STE7 (mitogen-activated protein kinase kinase), and FUS3/KSS1 (mitogen-activated protein kinase) in response to signals from both Cdc42 and the heterotrimeric G proteins associated with transmembrane pheromone receptors. ACF5 is a human cDNA encoding a protein kinase homologous to STE20. This protein kinase, also designated HPK/GCK-like kinase (HGK), has nucleotide sequences that encode an open reading frame of 1165 amino acids with 11 kinase subdomains. HGK is a serine/threonine protein kinase that specifically activated the c-Jun N-terminal kinase (JNK) signaling pathway when transfected into 293T cells, but does not stimulate either the extracellular signal-regulated kinase or p38 kinase pathway. HGK also increased AP-1-mediated transcriptional activity in vivo. HGK may be a novel activator of the JNK pathway. The cascade may look like this: HGK -> TAK1 -> MKK4, MKK7 -> JNK kinase cascade, which may mediate the TNF-alpha signaling pathway.

Paragraph beginning at line 4 of page 161 has been amended as follows:

ACF8 protein sequence (SEQ ID NO:127):

Gene name: Phospholipase A2, group IVC (cytosolic, calcium-independent)

Unigene number: Hs.18858

Probeset Accession #: AA054087

Protein Accession #: NP_003697.1

Pfam: none identified

Summary: ACF8 is a membrane-bound, calcium-independent PLA2, named cPLA2-gamma. The sequence encodes a 541-amino acid protein containing a domain with significant homology to the catalytic domain of the 85-kDa cPLA2 (cPLA2-alpha). cPLA2-gamma does not contain the regulatory calcium-dependent lipid binding (CaLB) domain found in cPLA2-alpha. cPLA2-gamma does contain two consensus motifs for lipid modification, a prenylation motif (-CCLA) at the C terminus and a myristoylation site at the N terminus. cPLA2-gamma demonstrates a preference for arachidonic acid at the sn-2 position of phosphatidylcholine as compared with palmitic acid. cPLA2-gamma encodes a 3-kilobase message, which is highly expressed in heart and skeletal muscle, suggesting a specific role in these tissues.

10021660-120601

Paragraph beginning at line 34 of page 161 has been amended as follows:

ACG1 protein sequence (SEQ ID NO:128):

Gene name: Carbohydrate (chondroitin 6/keratan) sulfotransferase 1

Unigene number: Hs.104576

Probeset Accession #: AA868063

Protein Accession #: NP_003645.1

Pfam: none identified

Summary: Chondroitin 6-sulfotransferase (C6ST) is the key enzyme in the biosynthesis of chondroitin 6-sulfate, a glycosaminoglycan implicated in chondrogenesis, neoplasia, atherosclerosis, and other processes. C6ST catalyzes the transfer of sulfate from 3'-phosphoadenosine 5'-phosphosulfate to carbon 6 of the N-acetylgalactosamine residues of chondroitin.

Paragraph beginning at line 55 of page 161 has been amended as follows:

ACG5 protein sequence (SEQ ID NO:129):

Gene name: Multimerin

Unigene number: Hs.268107

Probeset Accession #: U27109

Protein Accession #: AAC52065

Signal sequence: prediction underlined

Pfam: EGF-like domain, C1q domain

Summary: Multimerin is a massive, soluble protein found in platelets and in the endothelium of blood vessels. Multimerin is composed of varying sized, disulfide-linked multimers, the smallest of which is a homotrimer. Multimerin is a factor V/Va-binding protein and may function as a carrier protein for platelet factor V. Northern analyses show a 4.7-kilobase transcript in cultured endothelial cells, a megakaryocytic cell line, platelets, and highly vascular tissues. The multimerin cDNA can encode a protein of 1228 amino acids with the probable signal peptide cleavage site between amino acids 19 and 20. The protein is predicted to be hydrophilic and to contain 23 N-glycosylation sites. The adhesive motif RGDS (Arg-Gly-Asp-Ser) and an epidermal growth factor-like domain were identified. Multimerin contains a probable coiled-coil structures in the central portion of its sequence. Additionally, the carboxyl-terminal region of multimerin resembles the globular, non-collagen-like, carboxyl-terminal domains of several other trimeric proteins, including complement C1q and collagens type VIII and X.

Paragraph beginning at line 32 of page 162 has been amended as follows:

ACC6 protein sequence (SEQ ID NO:131):

Gene name: Homo sapiens cDNA FLJ11502 fis, clone HEMBA1002102, weakly similar to ANKRYIN

Unigene number: Hs.213194

Probeset Accession #: AA187101

Protein Accession #: none

Pfam: ankyrin repeats

10021660-120601

Paragraph beginning at line 45 of page 162 has been amended as follows:

ACC7 protein sequence (SEQ ID NO:132):

Gene name: Human RAL A gene

Unigene number: Hs.6906

Probeset Accession #: AA083572 cluster

Protein Accession #: P11233

Pfam: ras

Features: CAAX motif is underlined

Summary: The RALA gene encodes a low molecular mass ras-like GTP-binding protein that shares about 50% similarity with the ras proteins. GTP-binding proteins mediate the transmembrane signaling initiated by the occupancy of certain cell surface receptors. The RALA gene maps to 7p22-p15.

Paragraph beginning at line 63 of page 162 has been amended as follows:

ACC9 protein sequence (SEQ ID NO:133):

Gene name: KIAA0955 protein

Unigene number: Hs.10031

Probeset Accession #: AA027168

Protein Accession #: BAA76799.1

Pfam: CARD (Caspase recruitment domain)

Summary: Gene was originally isolated as a brain cDNA. The coding region contains a CARD domain, suggesting involvement in apoptotic signaling pathways.

Paragraph beginning at line 14 of page 163 has been amended as follows:

ACF6 Protein sequence (SEQ ID NO:134):

Gene name: Homo sapiens cDNA FLJ10669 fis, clone NT2RP2006275, weakly similar to Microtubule-associated protein 1B [CONTAINS: LIGHT CHAIN LC1]

Unigene number: Hs.66048

Probeset Accession #: AA609717

Protein Accession #: BAA91743.1

Pfam: none identified

Summary: The cDNA for FLJ10669 was originally isolated from NT2 neuronal precursor cells (teratocarcinoma cell line) after 2-weeks of retinoic acid (RA) treatment. The protein sequence has similarity to microtubule-associated protein 1B (MAP-1B), suggesting a function for ACF6 in the regulating the cytoskeleton.